



ENVIRONMENTAL
HEALTH
PERSPECTIVES

<http://www.ehponline.org>

Residential Proximity to Methyl Bromide Use and Birth Outcomes in an Agricultural Population in California

**Alison Gemmill, Robert B. Gunier, Asa Bradman,
Brenda Eskenazi and Kim G. Harley**

<http://dx.doi.org/10.1289/ehp.1205682>

Online 19 April 2013

Residential Proximity to Methyl Bromide Use and Birth Outcomes in an Agricultural Population in California

Alison Gemmill¹, Robert B. Gunier², Asa Bradman², Brenda Eskenazi², Kim G. Harley²

¹ Department of Demography, University of California, Berkeley, Berkeley, California, USA

² Center for Environmental Research and Children's Health (CERCH), School of Public Health, University of California, Berkeley, Berkeley, California, USA

Corresponding Author:

Kim G. Harley, Ph.D.

Center for Environmental Research and Children's Health (CERCH)

School of Public Health

University of California, Berkeley

1995 University Ave., Suite 265

Berkeley, CA 94602

Email: kharley@berkeley.edu

Phone: (510) 643-1310

FAX: (510) 642-9083.

Running title: Methyl bromide and birth outcomes

Keywords: birth outcomes, birth weight, fumigants, methyl bromide, pesticides, residential proximity

Acknowledgements: This publication was made possible by research supported by the following grants: PO1 ES009605 from NIEHS, RD83451301 from EPA, and HD07275 from

NICHD. We also acknowledge the CHAMACOS participants and families, without whom this - study would not have been possible. -

Competing interests declaration: The authors declare they have no actual or potential - competing financial interests. -

Abbreviations:

BMI	Body Mass Index
CHAMACOS	Center for the Health Assessment of Mothers and Children of Salinas
CI	Confidence Interval
GPS	Global Positioning System
GIS	Global Information System
MeBr	Methyl Bromide
PLSS	Public Land Survey System
PUR	Pesticide Use Reporting

Abstract

Background: Methyl bromide, a fungicide often used in strawberry cultivation, is of concern for residents who live near agricultural applications because of its toxicity and potential for drift. Little is known about the effects of methyl bromide exposure during pregnancy.

Objective: We investigated the relationship between residential proximity to methyl bromide use and birth outcomes.

Methods: Participants were from the CHAMACOS study (n=442), a longitudinal cohort study examining the health effects of environmental exposures on pregnant women and their children in an agricultural community in Northern California. Using data from the California Pesticide Use Reporting system, we employed a geographic information system (GIS) to estimate the amount of methyl bromide applied within 5 km of a woman's residence during pregnancy. Multiple linear regression models were used to estimate associations between trimester-specific proximity to use and birth weight, length, head circumference, and gestational age.

Results: High methyl bromide use (vs. no use) within 5 km of the home during the second trimester was negatively associated with birth weight ($\beta = -113.1$ g; CI: -218.1, -8.1), birth length ($\beta = -0.85$ cm; CI: -1.44, -0.27), and head circumference ($\beta = -0.33$ cm; CI: -0.67, 0.01). These outcomes were also associated with moderate methyl bromide use during the second trimester. Negative associations with fetal growth parameters were stronger when larger (5 km and 8 km) versus smaller (1 km and 3 km) buffer zones were used to estimate exposure.

Conclusions: Residential proximity to methyl bromide use during the second trimester was associated with markers of restricted fetal growth in our study.

Introduction

Methyl bromide is a fumigant used for pre-plant soil preparation. Because of concerns about its effect on the ozone layer, methyl bromide was banned in 2005 under the Montreal Protocol (United Nations Environment Programme 2000). Before the ban, methyl bromide was one of the most heavily used pesticides in the state of California, with nearly 5 million kg applied in 2000 (California Department of Pesticide Regulation (DPR) 2001b). Although methyl bromide use has declined in recent years, over 1.75 million kg were applied throughout the state in 2010 due to critical use exemptions for strawberries and other agricultural commodities (California DPR 2012c). Critical use of methyl bromide in the United States has declined markedly since 2005, but authorizations for use are still granted by the Environmental Protection Agency (United States Government 2012).

Animal studies of methyl bromide exposure suggest the potential for developmental toxicity (Subcommittee for the Review of the Risk Assessment of Methyl Bromide 2000). In an inhalation study of New Zealand white rabbits, exposure to 80-ppm during gestational days 7 to 19 resulted in decreased fetal body weights and increased rates of fetal malformations including gall bladder agenesis and fused sternbrae (Breslin et al. 1990). In another inhalation study, pups born to dams exposed to 20- and 70-ppm during gestation had increased rates of skull ossification defects (Sikov et al. 1981). A similar study of rats found dose-dependent reductions in fetal body and brain weights, as well as reduced cerebral cortex widths (American Biogenics Corporation 1986). Based on this limited evidence, the California Department of Pesticide Regulation lists methyl bromide as a probable developmental toxicant (Subcommittee for the Review of the Risk Assessment of Methyl Bromide 2000).

Several human poisoning case studies indicate that high levels of acute methyl bromide exposure are associated with dermal burns, neurological impairment, and death (Breeman 2009; Langård et al. 1996; Lifshitz and Gavrilov 2000). Much less is known about the effects of chronic, low-level exposure in human populations (Subcommittee for the Review of the Risk Assessment of Methyl Bromide 2000). In a sample of 56 male workers, long-term occupational exposure to methyl bromide was associated with chronic symptoms of dizziness, numbness, nightmares, and fatigue (Kishi et al. 1991). Other studies have investigated the health effects of residential proximity to methyl bromide use as part of larger analyses of ambient pesticide exposure. To date, one study reported evidence of an association with prostate cancer (Cockburn et al. 2011), but studies of breast cancer (Reynolds et al. 2004), childhood cancers (Reynolds et al. 2005), and autism spectrum disorders (Roberts et al. 2007) did not report associations with proximity to methyl bromide use.

Unlike most pesticides, fumigants such as methyl bromide have a high vapor pressure and are more likely to drift off-site (Woodrow and Krieger 2007). It is estimated that 30-50% of agricultural applications of methyl bromide are released into the air even when protective measures such as plastic tarps are in place (Honaganahalli and Seiber 2000; Majewski et al. 1995; Yagi et al. 1993). Because inhalation is the primary route of exposure to methyl bromide, ambient air concentrations are important for assessing exposure to nearby residents (Subcommittee for the Review of the Risk Assessment of Methyl Bromide 2000). In agricultural areas of California, measured outdoor concentrations of methyl bromide averaged 0.1 – 7.7 ppb over an 8-week monitoring period in 2000 (California Air Resources Board 2001), or up to one-tenth of the levels associated with adverse outcomes in some animal studies (Breslin et al. 1990;

Sikov et al. 1981). In addition, methyl bromide has been detected in air over 70 km away from the nearest application site (Honaganahalli and Seiber 2000).

Chronic exposure to methyl bromide is difficult to assess because there are no exposure biomarkers (Minnesota Department of Health 1999). The few studies that have assessed chronic methyl bromide exposure have done so by residential proxy or occupational history reports (Cockburn et al. 2011; Kishi et al. 1991; Reynolds et al. 2004, 2005; Roberts et al. 2007). Studies in California can utilize extensive information on the timing, location, and amount of agricultural methyl bromide applications, which must be reported to the state's Pesticide Use Reporting (PUR) system (California DPR 2012b).

In the current study we examine whether residential proximity to methyl bromide applications is associated with fetal growth and gestational length in a cohort of pregnant women living in an agricultural community in the Salinas Valley of California. We linked California PUR data with birth outcome and demographic data for pregnant women participating in the Center for the Health Assessment of Mothers and Children of Salinas (CHAMACOS), a longitudinal cohort study examining the health effects of environmental exposures on pregnant women and their children living in an agricultural community in Northern California.

Methods

Participants: Women were recruited for the CHAMACOS study if they presented for prenatal care at one of six partnering community clinics located throughout the Salinas Valley between October 1999 and October 2000. Women were eligible for the study if they were at least 18 years of age, ≤ 20 weeks gestation, English or Spanish speaking, Medi-Cal eligible, and planning to deliver at the county hospital.

A total of 601 women were enrolled, and after losses due to miscarriage and loss-to-follow-up, birth weight information was available for 536 children. Women were excluded from the present analysis if they delivered twins ($n=5$), stillbirths ($n=3$), or infants born at less than 500g ($n=1$), as well as women who were diagnosed with pregnancy-induced hypertension ($n=15$) or diabetes ($n=26$). Women were also excluded if residential proximity to methyl bromide use could not be assessed for at least 75 days during one trimester of pregnancy ($n=44$). Participants who were excluded were likely to have moved at least once during pregnancy (any residential moves: 85% vs. 41%) and tended to be less educated (high school graduate: 15% vs. 22%) and lower parity (nulliparous: 46% vs. 31%) than participants included in the present analysis. The final sample size was 442 mothers with residential information available for at least one trimester of pregnancy. Written informed consent was provided by all women, and the study was approved by the Committee for the Protection of Human Subjects at UC Berkeley.

Data Collection: Women were interviewed at the time of enrollment (median: 13 weeks gestation), at the end of the second trimester of pregnancy (median: 26 weeks gestation), and following delivery. All interviews were conducted by bilingual interviewers using structured questionnaires in English or Spanish. Interviews obtained demographic information including maternal age, family income, and number of years lived in the United States, as well as BMI (based on self-reported pre-pregnancy weight and measured height), and pregnancy history. Information on alcohol, tobacco, drug, and caffeine use was also collected. Women were asked about their occupational status during pregnancy, including work in agriculture. Residential history information was collected by asking participants if they had moved since the last interview, and, if so, the dates of all moves.

A home inspection was conducted shortly after enrollment (median: 16 weeks gestation) and when the child was six months old. For both visits, latitude and longitude coordinates of the participant's home were determined using a handheld global positioning system (GPS) unit. Residential mobility during pregnancy was common, with 53% of all participants moving at least once during pregnancy. For this analysis, a woman was included in the sample if her residential location (latitude and longitude coordinates) was known for 75 days or more of at least one trimester of pregnancy (trimester 1: n=338; trimester 2: n=408; trimester 3: n=390). Since we collected latitude and longitude coordinates at two time points, both residential locations were known for women who moved once during pregnancy. For women who moved more than once, residential location was often known for at least one trimester, allowing us to include the woman in the analyses of that trimester and exclude her from the analyses of trimesters where her residence was unknown.

Birth outcome data were collected from medical records and abstracted by a registered nurse. Delivery data included information on pregnancy complications, maternal weight at delivery, infant birth weight in grams, length, head circumference, and gestational age at birth.

Geographic-Based Estimates of Methyl Bromide Use: We estimated agricultural use of methyl bromide near each woman's residence using the following variables from the 1999-2000 PUR data: pounds of active ingredient applied, application date, and location. The location of pesticide application is reported for each square mile section (approximately 1.6 km by 1.6 km) defined by the Public Land Survey System (Figure 1). Prior to analysis, the PUR data were edited to correct for likely outliers with unusually high application rates using previously developed methods (Gunier 2001).

For each woman, we calculated the amount of methyl bromide applied within a 5 km radius around her home using a Geographic Information System (GIS). Detailed descriptions of the equations and methods that were used to calculate pesticide use have been published previously (Gunier et al. 2011; Nuckols et al. 2007). Briefly, we calculated nearby methyl bromide by summing the kilograms applied in all 1.6 km x 1.6 km Public Land Survey System (PLSS) sections that fell within 5 km of the maternal residence (Figure 2). For sections that intersected the 5 km buffer, we weighted the amount of methyl bromide applied in that section by the proportion of land area that was included in the buffer. We summed these totals over each day of a trimester interval, yielding an estimate of the total amount of methyl bromide (in kilograms) applied within 5 km of the maternal residence during each trimester of pregnancy.

We selected a 5 km buffer distance for this analysis because it best captures the spatial scale most strongly correlated with measured ambient methyl bromide concentrations (Honaganahalli and Seiber 2000; Li et al. 2005; Ross et al. 1996). In particular, Li et al. (2005) investigated the empirical relationship between methyl bromide use and ambient air concentrations in the same agricultural region (Salinas Valley) and time period of interest (2000) as our study using a variety of spatiotemporal scales. They found that parameters for the best-fit model of ambient methyl bromide concentrations at monitoring sites were average methyl bromide use during a 7-8 week period over the 7 x 7 mile area surrounding the monitoring site ($R^2 = 0.95$). Accordingly, we chose a GIS buffer radius of 5 km (3.1 miles) around a residence to adequately capture this 7 x 7 mile area.

We analyzed trimester-specific sums of methyl bromide use in kilograms for each woman as both continuous and categorical variables. We log-transformed (\log_{10}) continuous methyl bromide variables to reduce the influence of outliers and improve the linear fit of the model. In

the categorical analyses, we compared the baseline category of no use (< 1 kg) with moderate use (exposure $<$ median) and high use (exposure \geq median) groups. To create categories of moderate and high use that were consistent across each trimester, the cut-point used was the average of the median values in each of the three trimesters.

Statistical Analysis: We used linear regression models to estimate associations between methyl bromide usage during each trimester of pregnancy and birth weight, length, head circumference, and length of gestation. All models of birth weight, length, and head circumference were adjusted for gestational age and gestational age squared to estimate the association with fetal growth while adjusting for length of gestation. Potential confounders considered *a priori* were maternal age and week of initiating prenatal care as continuous variables, and parity (nulliparous vs. multiparous), infant sex, mother's country of birth (born in Mexico vs. other), pre-pregnancy body mass index (BMI), family income (at or below vs. above the federal poverty threshold) and mother's work status during pregnancy (ever paid work vs. no paid work) as categorical variables. Missing covariate values for income level (n=22) were substituted with information from the closest follow-up interview. For women missing height (n=8), pre-pregnancy BMI was predicted from a regression model that included age, weight, alcohol consumption, and education as predictor variables. We included covariates in the model if their exclusion changed the estimated coefficient for the main effect by more than 10% or if they predicted the outcome ($p < 0.10$). Smoking and illicit drug use were not included in the model because few women reported these behaviors and they did not influence main effect estimates. Adjusting for exposure to environmental tobacco smoke and alcohol and caffeine use during pregnancy also did not affect the coefficients of interest.

We conducted sensitivity analyses of alternative buffer distances around the maternal residence, including two smaller distances (1 km and 3 km) and a buffer distance that is slightly larger (8 km) than the 7 x 7 square mile area shown to be optimal for estimating exposure in the Li et al. study (2005). For these analyses, methyl bromide use near the residence was categorized into a dichotomous variable [any use (≥ 1 kg) vs. no use (<1 kg)]. We also carried out additional sensitivity analyses to investigate the influence of outliers in both the dependent and independent variables.

We conducted all statistical analyses using STATA version 11 statistical software (StataCorp LP, College Station, Texas).

Results

Most of the women in the study were Latina (96%) and born in Mexico (85%) (Table 1). About half were recent immigrants to the United States, having lived in the country for five years or less. Most mothers lived in families with incomes less than 200% of the federal poverty level (96%) and 78% had not graduated from high school. The median age was 25 (SD: 5), and few women smoked during pregnancy (6%); consumption of ≥ 1 alcoholic drink per week (1%) and illegal drug use (2%) were also low (data not shown). Over half of the sample was either overweight (40%) or obese (22%). Approximately 4% of deliveries were born with low birth weight (<2500 g) and 7% were preterm (<37 weeks gestational age). Characteristics of women in each trimester sub-sample were similar to those of the total sample (see Supplemental Material, Table S1).

Table 2 shows the distribution of methyl bromide use for each trimester for the buffer distance of interest (5 km) and the three additional buffer distances used in sensitivity analyses (1, 3, and 8

km). The range of methyl bromide use near residences was similar for trimesters 2 and 3 for each buffer distance, while values for trimester 1 were consistently larger due to the presence of 12 outliers (defined as $> Q3 + (1.5 * IQR)$, where $Q3$ is the value of the 75th percentile and IQR is the interquartile range). Not surprisingly, the proportion of women living near methyl bromide applications increased as the buffer distance increased. For example, 85% of women lived within 8 km of some amount of methyl bromide use during the first trimester compared to 78%, 60% and 16% of women for the 5 km, 3 km, and 1 km distances respectively. For the primary buffer distance used in this analysis (5 km), the estimated median amounts of methyl bromide use ranged from 6,010 to 8,201 kg, and the 95th percentile values ranged from 53,685 to 68,893 kg.

Table 3 presents estimated associations between methyl bromide use within 5 km of the home as a continuous variable (log 10) and the four birth outcomes of interest. In minimally adjusted analyses, increases in methyl bromide use near residences in the second trimester were associated with decreases in birth weight, birth length, and head circumference. After adjusting for confounders, associations remained; each ten-fold increase of methyl bromide use in the second trimester was associated with a 21.4 g (CI: -43.2, 0.4) decrease in birth weight, a 0.16 cm (CI: -0.28, -0.04) decrease in birth length, and a 0.08 cm (CI: -0.15, -0.01) decrease in head circumference. Model estimates did not support associations between methyl bromide use in the first or third trimesters and fetal growth. Residential proximity to methyl bromide use was also positively associated with gestational age in the first trimester in both crude and adjusted analysis (adjusted $\beta = 0.13$ week; CI: 0.03, 0.22).

Proximity to methyl bromide use was also analyzed as a three-level categorical variable (Table 4). The findings are similar to those from the continuous variable analysis. Using the 5 km buffer, moderate use of methyl bromide (vs. no use) nearby during the first trimester was

associated with a 0.70 week increase in gestational age (CI: 0.21, 1.18) and high use was associated with a 0.59 week increase (CI: 0.12, 1.06). Moderate and high methyl bromide use within 5km of the home during the second trimester were negatively associated with birth weight ($\beta = -93.1$ g; CI: -198.0, 11.7 and $\beta = -113.1$ g; CI: -218.1, -8.1, respectively) and birth length ($\beta = -0.62$ cm; CI: -1.20, -0.03 and $\beta = -0.85$ cm; CI: -1.44, -0.27, respectively), with stronger associations with high versus moderate use. In addition, proximity to moderate or high use in the second trimester was negatively associated with head circumference ($\beta = -0.42$ cm; CI: -0.76, -0.08 and $\beta = -0.33$ cm; CI: -0.67, 0.01, respectively). There was little evidence of associations between proximity to methyl bromide use during the first and third trimesters and birth weight, length, or head circumference.

Sensitivity analyses were conducted using different buffer distances around the maternal residence. Results for second trimester methyl bromide use (any vs. none) are presented in Table 5 using 1 km, 3 km, 5 km and 8 km distances. Proximity to methyl bromide use within 1 km and 3 km during the second trimester was negatively associated with body length. Additionally, the 3 km distance yields marginally significant associations with reduced birth weight ($p=0.06$), reduced head circumference ($p=0.08$), and increased gestational age ($p=0.05$). Unlike the smaller buffer distances, the 8 km analysis was more consistent with the 5 km buffer analysis; proximity to methyl bromide use during the second trimester was negatively associated with birth weight, length and head circumference, and associations were stronger than estimates from the 5 km analysis. Sensitivity analyses with other trimesters yielded few statistically significant associations: in the 3 km analysis there was a significant positive association with first trimester exposure and gestational age (see Supplemental Material, Table S2).

Discussion

Living within 5 km of methyl bromide use in the second trimester of pregnancy was associated with decreased birth weight, length, and head circumference, despite a positive association between gestational age and residential proximity to methyl bromide use in the first trimester. The estimates for birth weight were only marginally significant ($p=0.08$) among those women living near moderate methyl bromide use, while those for head circumference were marginally significant ($p=0.06$) among those with high use. Model estimates did not support associations between exposure to methyl bromide during the first and third trimester and fetal growth parameters. Associations with nearby methyl bromide use analyzed as a continuous variable were consistent with those based on categorical exposures: each 10-fold increase in kilograms of methyl bromide applied within 5 km of the home during the second trimester of pregnancy was associated with a 24.1 g decrease in mean birth weight, a 0.16 cm decrease in mean birth length, and a 0.08 cm decrease in mean head circumference.

The estimated decrease in mean birth weight associated with living near high use during the second trimester (113g) was about half of the 250 g birth weight decrease generally associated with maternal smoking (Kramer 1987). Nevertheless, the findings suggest that the second trimester may be a critical period for effects of exposure for gestational growth. Findings from studies investigating the effects of maternal smoking (Reeves and Bernstein 2008) and prenatal exposure to wildfires (Holstius et al. 2012) suggest that both the 2nd and 3rd trimesters may be critical periods of growth. In these studies, the authors cite oxidative stress, hypoxia, and suppression of placenta growth hormone as plausible mechanisms.

The results from our sensitivity analyses suggested stronger associations as the buffer distance increased, with negative associations with all three fetal growth parameters for usage within 5 km and 8 km buffer areas. The lack of consistent associations with shorter buffer distances may have been at least partly due to exposure misclassification, as Li et al. (2005) concluded that using smaller buffers (<5 km) to determine methyl bromide use underestimates measured outdoor air concentrations. Previous studies using GIS to assess residential exposure to methyl bromide have used a much smaller buffer distance than the present analysis (Cockburn et al. 2011; Reynolds et al. 2004; Rull et al. 2009). These studies selected a buffer distance of 500 m or 0.5 mile (~ 800 m) for all pesticides, but did not use air monitoring data to confirm the validity of these buffer sizes for exposure assessment. In Cockburn et al. (2011), a 500 m buffer was justified based on a previous study that observed high specificity for serum DDE concentrations using a 1 km GIS buffer to estimate organochlorine exposure (Ritz and Costello 2006). While this shorter buffer distance might be appropriate for assessing exposure to other pesticides, it does not appear to be sufficient for assessing exposure to fumigants, which are likely to drift much further away from the application site (Honaganahalli and Seiber 2000; Li et al. 2005; Woodrow and Krieger 2007). Regulations in California recognize the potential for methyl bromide to drift far from the application site and restrict methyl bromide use at the township level (6 x 6 square mile sections) to a maximum of 77,848 kg in a calendar month to reduce outdoor air concentrations (California DPR 2010).

The main limitations of this study are that exposure to methyl bromide was based on residential proximity to reported agricultural use, not measured personal air concentrations. Thus time spent in areas with no methyl bromide use or exposure away from home, including at work, was not taken into account. In addition, in our study area and time period, use of chloropicrin, a

chemical often used in conjunction with methyl bromide applications, was very highly correlated with methyl bromide use (Spearman correlation=0.95). As a result, it is not possible to separate associations with methyl bromide from associations with chloropicrin in this study. Use of another pesticide, diazinon, was also correlated with methyl bromide use, but much less strongly (Spearman correlation=0.44). Future analyses will examine whether diazinon is associated with birth outcomes in this population.

Although we estimated exposure to a specific pesticide that was selected *a priori* based on toxicity and volume of use, other agricultural pesticides could also be associated with fetal growth. However, previous analyses of this cohort did not indicate associations between decreased fetal growth and organophosphate metabolites measured in maternal urine (Eskenazi et al. 2004) or organochlorine concentrations in maternal serum (Fenster et al. 2006).

Methyl bromide use is seasonal, with the most use occurring between August and October. Thus, it is possible that the association we found reflects seasonal patterns in birth weight rather than 2nd trimester methyl bromide use. However, we did not observe any seasonal patterns in birth weight in this population. Additionally, controlling for birth between February-April (corresponding to a second trimester during August-October) in the multivariable models did not change the results (data not shown).

We did not use a dispersion model incorporating meteorological data to estimate methyl bromide concentrations because dispersion models have not predicted fumigant concentrations as well ($R^2=0.55-0.82$) as the regression model approach ($R^2=0.95$) that we used (Honaganahalli and Seiber 2000; Li et al. 2005). Additionally, location of methyl bromide use was reported to one-square-mile units. The PUR reporting system does not contain information on applications to

specific fields, so it is not possible to ascertain where in the PLSS section the material was applied. However, the potential impact of this misclassification would be small because most PLSS sections fell entirely within the larger buffer distances (e.g. 5 and 8 km) used in our analysis.

Another potential limitation is that women who lived near high use of methyl bromide (vs. moderate or no use) were more likely to live in and around the city of Salinas, CA. It is possible that unmeasured factors related to this group of women might have confounded our results; however, associations remained after controlling for several potential confounders, including maternal education and income. In addition, associations with fetal growth parameters were seen with both moderate and high use of methyl bromide.

Previous studies of perinatal outcomes and agricultural pesticide exposures have relied on the address given in birth certificates and have not incorporated environmental monitoring information in the exposure models (Shirangi et al. 2011). In contrast, for most women, we were able to capture residential location throughout pregnancy, incorporate residential mobility, and estimate exposure separately for each trimester (Canfield et al. 2006). With slightly over half of our original sample moving at least once during pregnancy, accounting for residential history was important to reduce possible exposure misclassification. Furthermore, residential location was determined using a GPS, which is more accurate than geocoding self-reported addresses (Ward et al. 2005). Another strength of the study was that fetal growth parameters were obtained from medical records, which generally provide more accurate information than birth certificates (Lain et al. 2012). Additionally, because methyl bromide is highly volatile (USEPA 2006), very little exposure to methyl bromide occurs from dietary ingestion or residues brought home on the

clothes and skin of occupationally-exposed household members; therefore, exposure misclassification related to these sources is not a concern.

The results of this study suggest that living near agricultural applications of methyl bromide may be associated with decreased infant birth weight and other measures of fetal growth. Our finding that residential proximity to methyl bromide use in the first trimester of pregnancy was associated with longer duration of gestation is somewhat puzzling and counter to our expectations. No other epidemiologic studies have specifically examined methyl bromide exposure and fetal growth; therefore, additional studies are needed to confirm the associations we observed. Methyl bromide use in the Salinas Valley declined from 850,000 kg in 2000 to 565,000 kg in 2010 but still represented 29% of soil fumigant use (California DPR 2001a, 2012a). While critical use exemptions for methyl bromide use are being phased out (US Government 2012), the lack of fumigant alternatives may prolong its use in the state of California.

References

- American Biogenics Corporation. 1986. Two-generation reproduction study via inhalation in albino rats using methyl bromide. Study 450– 1525. Submitted to Methyl Bromide Panel, c/o Great Lakes Chemical Corporation, W.Lafayette, IN. Decatur, IL:American Biogenics Corporation.
- Breeman W. 2009. Methylbromide intoxication: a clinical case study. *Adv Emerg Nurs J* 31:153-160.
- Breslin WJ, Zabloutny CL, Bradley GJ, Lomax LG. 1990. Methyl bromide inhalation teratology study in New Zealand white rabbits. Final Report. K-000681–033. Study prepared by The Toxicology Research Laboratory, The Dow Chemical Company, Midland, MI for The Methyl Bromide Industry Panel, Chemical Manufacturers Association, Washington, DC.
- Canfield MA, Ramadhani TA, Langlois PH, Waller DK. 2006. Residential mobility patterns and exposure misclassification in epidemiologic studies of birth defects. *J Expo Sci Environ Epidemiol* 16:538-543.
- California Air Resources Board. 2001. Ambient Air Monitoring for Methyl Bromide and 1,3-Dichloropropene in Monterey and Santa Cruz Counties –Fall 2000. Sacramento, CA:California Air Resources Board. Available: <http://www.cdpr.ca.gov/docs/emon/pubs/tac/methylbr.htm> [accessed 27 September 2011].
- California DPR. 2001a. 2000 Annual Pesticide Use Report Preliminary Data: Monterey County Indexed by Chemical. Sacramento, CA:California DPR. Available: <http://www.cdpr.ca.gov/docs/pur/pur00rep/chemcnty/monter00.pdf> [accessed 20 June 2012].
- California Department of Pesticide Regulation (DPR). 2001b. Summary of Pesticide Use Report Data 2000, Indexed by Chemical: Preliminary Data. Sacramento, CA:California DPR. Available: <http://www.cdpr.ca.gov/docs/pur/pur00rep/00chem.htm> [accessed 27 March 2011].
- California DPR. 2010. Methyl Bromide Field Fumigation Regulations 10-002. Sacramento, CA:California DPR. Available: http://www.cdpr.ca.gov/docs/legbills/rulepkgs/10-002/text_final.pdf [accessed 27 September 2011].

- California DPR. 2012a. 2010 Annual Pesticide Use Report Indexed by Chemical, Monterey County. Sacramento, CA:California DPR. Available: http://www.cdpr.ca.gov/docs/pur/pur10rep/chemcnty/monter10_ai.pdf [accessed 6 June 2012].
- California DPR. 2012b. California Pesticide Use Reporting Program. Sacramento, CA:California DPR. Available: <http://www.cdpr.ca.gov/docs/pur/purmain.htm> [accessed 6 June 2012].
- California DPR. 2012c. The top 100 pesticides used pounds of active ingredients statewide in 2010 (all sites combined). Available: http://www.cdpr.ca.gov/docs/pur/pur10rep/top_100_ais_lbs10.pdf [accessed 23 March 2012].
- Cockburn M, Mills P, Zhang X, Zadnick J, Goldberg D, Ritz B. 2011. Prostate cancer and ambient pesticide exposure in agriculturally intensive areas in California. *Am J Epidemiol* 173:1280-1288.
- Eskenazi B, Harley K, Bradman A, Weltzien E, Jewell NP, Barr DB, et al. 2004. Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 112:1116-1124.
- Fenster L, Eskenazi B, Anderson M, Bradman A, Harley K, Hernandez H, et al. 2006. Association of in utero organochlorine pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 114:597-602.
- Gunier RB, Harnly ME, Reynolds P, Hertz A, Von Behren J. 2001. Agricultural pesticide use in California: pesticide prioritization, use densities, and population distributions for a childhood cancer study. *Environ Health Perspect* 109:1071-1078.
- Gunier RB, Ward MH, Airola M, Bell EM, Colt J, Nishioka M, et al. 2011. Determinants of agricultural pesticide concentrations in carpet dust. *Environ Health Perspect* 119:970-976.
- Holstius DM, Reid CE, Jesdale BM, Morello-Frosch R. 2012. Birth weight following pregnancy during the 2003 Southern California wildfires. *Environ Health Perspect* 120:1340-1345.
- Honaganahalli PS, Seiber JN. 2000. Measured and predicted airshed concentrations of methyl bromide in an agricultural valley and applications to exposure assessment. *Atmosph Environ* 34:3511-3523.
- Kishi R, Itoh I, Ishizu S, Harabuchi I, Miyake H. 1991. Symptoms among workers with long-term exposure to methyl bromide. An epidemiological study. *Sangyo Igaku* 33:241-250.

- Kramer MS. 1987. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ* 65:663-737.
- Lain SJ, Hadfield RM, Raynes-Greenow CH, Ford JB, Mealing NM, Algert CS, et al. 2012. Quality of data in perinatal population health databases: a systematic review. *Med Care* 50:e7-e20.
- Langård S, Rognum T, Fløtterød O, Skaug V. 1996. Fatal accident resulting from methyl bromide poisoning after fumigation of a neighbouring house; leakage through sewage pipes. *J Appl Toxicol* 16:445-448.
- Li LY, Johnson B, Segawa R. 2005. Empirical relationship between use, area, and ambient air concentration of methyl bromide. *J Environ Qual* 34:420-428.
- Lifshitz M, Gavrilov V. 2000. Central nervous system toxicity and early peripheral neuropathy following dermal exposure to methyl bromide. *J Toxicol Clin Toxicol* 38:799-801.
- Majewski MS, McChesney MM, Woodrow JE, Prueger JH, Seiber JN. 1995. Aerodynamic measurements of methyl bromide volatilization from tarped and nontarped fields. *J Environ Qual* 24:742-752.
- Minnesota Department of Health. 1999. Health consultation: exposure to methyl bromide as a result of fumigation at ADM Milling Company. Minneapolis, MN:Minnesota Department of Health. Available: <http://www.health.state.mn.us/divs/eh/hazardous/sites/hennepin/adm0399.pdf> [accessed 6 April 2011].
- Nuckols JR, Gunier RB, Riggs P, Miller R, Reynolds P, Ward MG. 2007. Linkage of the California Pesticide Use Reporting Database with spatial land use data for exposure assessment. *Environ Health Perspect* 115:684-689.
- Reeves S, Bernstein I. 2008. Effects of maternal tobacco-smoke exposure on fetal growth and neonatal size. *Expert Rev Obstet Gynecol* 3:719-730.
- Reynolds P, Hurley SE, Goldberg DE, Yerabati S, Gunier RB, Hertz A, et al. 2004. Residential proximity to agricultural pesticide use and incidence of breast cancer in the California Teachers Study cohort. *Environ Res* 96:206-218.
- Reynolds P, Von Behren J, Gunier RB, Goldberg DE, Harnly M, Hertz A. 2005. Agricultural Pesticide Use and Childhood Cancer in California. *Epidemiology* 16:93-100.

- Ritz B, Costello S. 2006. Geographic model and biomarker-derived measures of pesticide exposure and Parkinson's disease. *Ann N Y Acad Sci* 1076:378-387.
- Roberts E, English PB, Grether JK, Windham GC, Somberg L, Wolff C. 2007. Maternal residence near agricultural pesticide applications and autism spectrum disorders among children in the California Central Valley. *Environ Health Perspect* 115:1482-1489.
- Ross LJ, Johnson B, Kim KD, Hsu J. 1996. Prediction of methyl bromide flux from area sources using the ISCST model. *J Environ Qual* 25:885-891.
- Rull RP, Gunier RB, Von Behren J, Hertz A, Crouse V, Buffler PA, et al. 2009. Residential proximity to agricultural pesticide applications and childhood acute lymphoblastic leukemia. *Environ Res* 109:891-899.
- Shirangi A, Nieuwenhuijsen M, Vienneau D, Holman CD. 2011. Living near agricultural pesticide applications and the risk of adverse reproductive outcomes: a review of the literature. *Paediatr Perinat Epidemiol* 25:172-191.
- Sikov MR, Cannon WC, Carr DB, Miller RA, Montgomery LF, Phelps DW. 1981. Teratologic assessment of butylene oxide, styrene oxide and methyl bromide. Contract no. 210-78-0025. Battelle, Pacific Northwest Lab. Submitted to the Division of Biomedical and Behavioral Science, National Institute for Occupational Safety and Health, U.S. Department of Health and Human Services, Cincinnati, OH.
- Subcommittee for the Review of the Risk Assessment of Methyl Bromide, Committee of Toxicology, Board on Environmental Studies and Toxicology, National Research Council. 2000. *Methyl Bromide Risk Characterization in California*. Washington, DC:National Academy Press.
- United Nations Environment Programme (UNEP). 2000. *The Montreal Protocol on Substances that Deplete the Ozone Layer*. Nairobi, Kenya:UNEP.
- United States Environmental Protection Agency (USEPA). 2006. Report on Food Quality Protection Act (FQPA) Tolerance Reassessment and Risk Management Decision (TRED) for Methyl Bromide, and Reregistration Eligibility Decision (RED) for Methyl Bromide's Commodity Uses. Report EPA 738-R-06-026. Available: http://www.epa.gov/opp00001/reregistration/REDs/methyl_bromide_tred.pdf [accessed 26 Feb 2013].

- United States Government. 2012. Protection of Stratospheric Ozone: The 2013 Critical Use Exemption From the Phaseout of Methyl Bromide. Federal Register. Volume 77. Number 241. Available: <http://www.gpo.gov/fdsys/pkg/FR-2012-12-14/pdf/2012-30225.pdf> [accessed 21 February 2013].
- Ward MH, Nuckols JR, Giglierano J, Bonner MR, Wolter C, Airola M, et al. 2005. Positional accuracy of two methods of geocoding. *Epidemiology* 16:542-547.
- Whyatt RM, Rauh V, Barr DB, Camann DE, Andrews HF, Garfinkel R, et al. 2004. Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 112:1125-1132.
- Wolff MS, Engel S, Berkowitz G, Teitelbaum S, Siskind J, Barr DB, et al. 2007. Prenatal pesticide and PCB exposures and birth outcomes. *Pediatr Res* 61:243-50.
- Woodrow JE, Krieger RI. 2007. Assessing exposure to agricultural fumigants in outdoor and indoor air environments. In: *Assessing Exposures and Reducing Risks to People from the Use of Pesticides* (Krieger RI, ed). 2007. Washington, DC:American Chemical Society, 70-86.
- Xiang H, Nuckols JR, Stallones L. 2000. A geographic information assessment of birth weight and crop production patterns around mother's residence. *Environ Res* 82:160-167.
- Yagi K, Williams J, Wang, NY, Cicerone RJ. 1993. Agricultural soil fumigation as a source of atmospheric methyl bromide. *Proc Natl Acad Sci* 90:8420-8423.

Table 1. Characteristics of study population, CHAMACOS study, Salinas Valley, CA, 1999-2000 (N=442).

Characteristics	N (%)
Maternal Age	
<20	40 (9.1%)
20 – 24	163 (36.9%)
25 – 29	144 (32.6%)
30 – 34	66 (14.9%)
≥35	29 (6.6%)
Parity	
0	139 (31.5%)
≥1	303 (68.6%)
Race/Ethnicity	
Latina	426 (96.4%)
Non-Latina, White	7 (1.6%)
Other	9 (2.0%)
Marital Status	
Married or living as married	356 (80.5%)
Unmarried	86 (19.5%)
Maternal education	
≤ 6th grade	188 (42.5%)
Some high school	158 (35.8%)
High school graduate	96 (21.7%)
Family income	
At or below poverty	272 (61.5%)
Poverty – 200%	154 (34.8%)
>200%	16 (3.6%)
Country of birth	
United States	59 (13.4%)
Mexico	374 (84.6%)
Other	9 (2.0%)
Years of residence in the U. S.	
≤ 5 years	230 (52.0%)
6 – 10 years	99 (22.4%)
11+ years	63 (14.3%)
Entire life	50 (11.3%)
Work status during pregnancy	
Did not work	172 (38.9%)
Some field or agricultural work	181 (41.0%)
Other work	89 (20.1%)
Prepregnancy body mass index (kg/m ²)	
Underweight (<18.5)	2 (0.5%)
Normal (18.5 – 24.9)	170 (38.5%)
Overweight (25 – 29.9)	175 (39.6%)

Characteristics	N (%)
Obese (>30)	95 (21.5%)
Smoked during pregnancy	
Yes	26 (5.9%)
No	416 (94.1%)
Any moves during pregnancy	
Yes	182 (41.2%)
No	260 (58.8%)
Low birth weight ^a infant	
Yes	17 (3.9%)
No	425 (96.2%)
Preterm ^b infant	
Yes	30 (6.8%)
No	412 (93.2%)

a Low birth weight is defined as <2,500 g.

b Preterm birth is defined as <37 weeks' gestation.

Table 2. Distribution of methyl bromide use near residences of pregnant women, CHAMACOS study, Salinas Valley, CA, 1999-2000 (N=442).

Group	N	No Nearby MeBr Use (%)	25th ^a	50th ^a	75th ^a	95th ^a
1km radius						
Trimester 1	338	83.7	0	0	0	1,669
Trimester 2	408	86.3	0	0	0	766
Trimester 3	390	86.4	0	0	0	615
3km radius						
Trimester 1	338	40.2	0	1,498	13,102	27,718
Trimester 2	408	40.2	0	734	7,379	23,455
Trimester 3	390	42.1	0	291	7,318	23,686
5km radius						
Trimester 1	338	21.6	82	8,201	34,838	68,893
Trimester 2	408	25.5	0	7,419	29,676	53,685
Trimester 3	390	25.4	0	6,010	26,688	59,983
8km radius						
Trimester 1	338	14.5	2,150	16,278	88,413	163,893
Trimester 2	408	18.9	352	11,984	69,200	129,464
Trimester 3	390	18.7	352	10,751	68,333	143,690

Abbreviations: MeBr Methyl Bromide

^a Percentile (in kg)

Table 3. Associations [β (95% CI)] of proximity to methyl bromide use within a 5-km radius (\log_{10}) during pregnancy with fetal growth parameters and length of gestation, CHAMACOS study, Salinas Valley, CA, 1999-2000 (N=442).

Trimester	Birth weight (g)			Length (cm)			Head circumference (cm)			Gestational age (weeks)		
			p-value			p-value			p-value			p-value
Minimally adjusted ^a												
Trimester 1	-5.0	(-30.7, 20.8)	0.71	-0.02	(-0.15, 0.12)	0.82	-0.01	(-0.09, 0.07)	0.85	0.12	(0.02, 0.21)	0.02
Trimester 2	-23.4	(-45.9, -0.9)	0.04	-0.16	(-0.28, -0.03)	0.01	-0.08	(-0.16, -0.01)	0.03	0.00	(-0.09, 0.09)	0.95
Trimester 3	-3.4	(-26.6, 19.8)	0.78	0.01	(-0.11, 0.14)	0.82	-0.06	(-0.13, 0.02)	0.15	0.05	(-0.05, 0.15)	0.30
Adjusted ^b												
Trimester 1	-11.1	(-36.4, 14.3)	0.39	-0.03	(-0.17, 0.11)	0.69	-0.03	(-0.11, 0.05)	0.43	0.13	(0.03, 0.22)	0.01
Trimester 2	-21.4	(-43.2, 0.4)	0.05	-0.16	(-0.28, -0.04)	0.01	-0.08	(-0.15, -0.01)	0.04	-0.00	(-0.09, 0.09)	1.00
Trimester 3	-0.8	(-23.5, 21.8)	0.94	0.01	(-0.12, 0.14)	0.90	-0.05	(-0.12, 0.03)	0.22	0.03	(-0.06, 0.13)	0.52

a All models of birth weight, length, and head circumference adjust for gestational age and gestational age squared.

b Adjusted for maternal age, parity, prepregnancy body mass index, poverty, country of birth, and week of initiating prenatal care.

Table 4. Categorical analysis of associations [β (95% CI)] of proximity to methyl bromide use within a 5-km radius during pregnancy with fetal growth parameters and length of gestation, CHAMACOS study, Salinas Valley, CA, 1999-2000. (N=442)

Group	Range of use (kg)	N	Birth weight (g) ^a	p-value	Length (cm) ^a	p-value	Head circumference (cm) ^a	p-value	Gestational age (weeks) ^b	p-value
Trimester 1										
None	<1	73	Ref		Ref		Ref		Ref	
Moderate ^c	1 - 14,690	121	-13.8 (-141.1, 113.6)	0.83	-0.16 (-0.85, 0.54)	0.66	-0.25 (-0.64, 0.15)	0.21	0.70 (0.21, 1.18)	0.01
High ^d	14,690 - 148,137	144	-27.7 (-149.9, 94.5)	0.66	-0.09 (-0.76, 0.58)	0.80	-0.06 (-0.44, 0.32)	0.75	0.59 (0.12, 1.06)	0.01
Trimester 2										
None	<1	104	Ref		Ref		Ref		Ref	
Moderate	1 - 14,690	152	-93.1 (-198.0, 11.7)	0.08	-0.62 (-1.20, -0.03)	0.04	-0.42 (-0.76, -0.08)	0.02	-0.03 (-0.47, 0.41)	0.89
High	14,690 - 76,306	152	-113.1 (-218.1, -8.1)	0.04	-0.85 (-1.44, -0.27)	<0.01	-0.33 (-0.67, 0.01)	0.06	-0.06 (-0.51, 0.38)	0.78
Trimester 3										
None	<1	99	Ref		Ref		Ref		Ref	
Moderate	1 - 14,690	155	14.0 (-93.1, 121.1)	0.80	-0.04 (-0.64, 0.56)	0.90	-0.27 (-0.62, 0.07)	0.12	0.22 (-0.23, 0.68)	0.33
High	14,690 - 77,079	136	-23.0 (-133.2, 87.2)	0.68	0.02 (-0.60, 0.64)	0.95	-0.26 (-0.62, 0.09)	0.15	0.18 (-0.29, 0.64)	0.46

a Adjusted for maternal age, parity, prepregnancy body mass index, poverty, country of birth, week of initiating prenatal care, gestational age, and gestational age squared.

b Adjusted for maternal age, parity, prepregnancy body mass index, poverty, country of birth, and week of initiating prenatal care.

c Moderate use is defined as less than the median (14,690 kg) among participants with any methyl bromide exposure.

d High use is defined as greater than the median (14,690 kg) among participants with any methyl bromide exposure.

Table 5. Categorical analysis of associations [β (95% CI)] of proximity to any methyl bromide use (vs. none) in the second trimester with fetal growth parameters and length of gestation, CHAMACOS study, Salinas Valley, CA, 1999-2000. (N=442)

Group	N	Birth weight (g) ^a		p-value	Length (cm) ^a		p-value	Head circumference (cm) ^a		p-value	Gestational age (weeks) ^b		p-value
1km radius													
None	352	Ref			Ref			Ref			Ref		
Any	56	-61.7	(-181.2, 57.8)	0.31	-0.70	(-1.37, -0.04)	0.04	0.07	(-0.32, 0.45)	0.73	-0.23	(-0.73, 0.27)	0.37
3km radius													
None	164	Ref			Ref			Ref			Ref		
Any	244	-81.2	(-165.3, 2.9)	0.06	-0.56	(-1.03, -0.09)	0.02	-0.25	(-0.52, 0.02)	0.08	0.35	(-0.00, 0.70)	0.05
5km radius													
None	104	Ref			Ref			Ref			Ref		
Any	304	-103.1	(-196.5, -9.6)	0.03	-0.73	(-1.25, -0.21)	0.01	-0.37	(-0.68, -0.07)	0.02	-0.05	(-0.44, 0.35)	0.81
8km radius													
None	77	Ref			Ref			Ref			Ref		
Any	331	-150.7	(-254.9, -46.6)	0.01	-1.03	(-1.61, -0.46)	<0.01	-0.42	(-0.76, -0.08)	0.02	-0.02	(-0.46, 0.42)	0.94


a Adjusted for maternal age, parity, prepregnancy body mass index, poverty, country of birth, week of initiating prenatal care, gestational age, and gestational age squared.

b Adjusted for maternal age, parity, prepregnancy body mass index, poverty, country of birth, and week of initiating prenatal care.

Figure Legends


Figure 1. Distribution of methyl bromide use (kg/year) from the California Pesticide Use Reporting system by Section of the Public Land Survey System grid in the Salinas Valley, 1999-2000.

Figure 2. Methods used to determine proximity to methyl bromide use from the California Pesticide Use Reporting system by Section of the Public Land Survey System grid illustrated for 3, 5, and 8 kilometer radius buffers around a residence (•).



Data Source: California Department of Pesticide Regulation 2012b.

Figure 1



Data Source: California Department of Pesticide Regulation 2012b.

Figure 2